Coupling QGIS to Analyse Mass, Classification, and Trends in Global Meteorite Landings

Yourfee Kennedy, Biosphere Reserves Management, MSc. Eberswalde University for Sustainable Development, Germany.

Abstract

This research report explores the integration of quantum geographic information systems (QGIS) with meteorite impact data to conduct a comprehensive analysis of the mass distribution, classification, and time trends of global meteorite impacts. This research aims to use geographic information system (GIS) technology to reveal hidden patterns and relationships in meteorite impact data to improve our understanding of meteorite dynamics and planetary processes. The approach involves collecting data from reputable sources such as the Meteorological Society database and the NASA Meteorite Research Database, and then preprocessing and analysing it using QGIS spatial analysis tools. The findings show regional and temporal changes in meteorite strikes, providing information on mass distribution, categorization, and temporal trends. The findings help to further planetary research and hazard assessment by revealing the frequency, distribution, and features of meteorite impacts worldwide.

Keywords: Meteorite impacts, QGIS, Geographic Information System, Spatial analysis, Temporal trends, Mass distribution, Classification, Planetary science, Hazard assessment, and Spatial variability.

1 Introduction

Meteorite impacts have long been a source of fascination for scientists and the general public alike, as they provide valuable insights into the composition and history of our solar system. To advance our understanding of celestial bodies and evaluate potential risks to Earth, it is vital to comprehend the spatial distribution, mass distribution, classification, and temporal trends of meteorite impacts. In recent times, Geographic Information Systems (GIS) have emerged as powerful tools for analysing spatial data, offering the ability to visualize, analyse, and interpret complex spatial patterns. In this investigation, we examine the integration of Quantum Geographic Information System (QGIS) with meteorite impact data to conduct a comprehensive analysis of mass distribution, classification, and temporal trends in global meteorite impacts. By harnessing the advanced spatial analysis capabilities of QGIS, we aim to uncover concealed patterns and relationships in meteorite impact data that can contribute to our understanding of meteorite dynamics and planetary processes (Xiao Long,2008).

2 Literature Review

Previous studies have investigated various aspects of meteorite landings using GIS techniques, focusing on spatial patterns, classification, and temporal trends. For instance, (Beckett J. R., et al. 2006) utilized GIS to analyze the spatial distribution of meteorite landings and identified hotspots of meteorite activity in certain geographic regions. Similarly, (Xia et al. 2016) conducted a classification analysis of meteorite samples using GIS-based techniques, revealing correlations between meteorite composition and geographic origin. These studies highlight the potential of GIS in enhancing our understanding of meteorite phenomena and their implications for planetary science. Additionally, advancements in GIS technology have enabled researchers to explore temporal trends in meteorite landings over time. (Kinoshita, K et al. 2016) employed temporal GIS techniques to analyze the temporal variability of meteorite landings and identified temporal clusters of meteorite activity. These findings underscore the importance of temporal analysis in elucidating the dynamic nature of meteorite landings and their underlying mechanisms Notwithstanding these developments, there is still a need for more research that combines GIS with meteorite landing data to do complete assessments of mass distribution, categorization, and time patterns. This project aims to close this gap by combining QGIS and meteorite landing data to provide new insights into the spatial and temporal dynamics of meteorite events.

3 Methodology

The methodology employed in this study consists of a sequence of phases, which include data collection, preprocessing, examination, and elucidation. Initially, information regarding meteorite impacts will be collected from credible sources like the Meteoritical Society Database and the NASA Meteorite Studies Database. This dataset will encompass various particulars, such as the meteorite's name, weight, categorization, descent, and the year of impact. Following this, the meteorite impact data will be imported into QGIS for preprocessing and analytical purposes. Preprocessing tasks may encompass activities such as data refinement, structuring, and spatial referencing to ensure smooth incorporation with the spatial analysis tools within QGIS. Spatial analysis techniques like spatial clustering, hotspot analysis, and interpolation will be utilized to explore spatial patterns and the distribution of meteorite impacts. An investigation into temporal patterns and variations in meteorite impacts will be conducted using temporal analysis methodologies. This procedure may involve the utilization of time-series analysis, trend analysis, and temporal clustering techniques to reveal trends and anomalies in the temporal occurrence of meteorite impacts. In conclusion, the findings of the investigation will be interpreted to provide insights into the distribution of mass, classification, and temporal trends in global meteorite impacts. These findings will be discussed in the context of existing literature and their implications for planetary science and risk assessment (M. K. Weisberg, et al. 2006).

4 Results

4.1 Heatmap of Meteorite Density by Fall Type



Figure 1. Shows the Density of Meteorite by fall type.

4.2 Temporal Trends in Meteorite Landings



Figure 2. Shows the temporal trends in Meteorite landings.

4.3 Spatial Distribution of Meteorite Classes



Figure 3. Shows the spatial distribution of meteorites per class.

4.4 Spatial Distribution of Meteorite Mass Worldwide



Figure 4. Shows the spatial distribution of meteorite mass.

5 Discussion

The analysis of meteorite impacts using GIS technology has provided valuable insights into various aspects of meteorite dynamics, including mass distribution, classification, and temporal trends. The results of this study shed light on the spatial and temporal patterns of meteorite impacts worldwide, contributing to our understanding of celestial phenomena and their implications for planetary science.

5.1 Mass Distribution and Spatial Patterns

Figure 1 shows the global meteorite impact density by fall type .The heat map shows significant spatial variation in meteorite density, with some areas having higher density of meteorite impacts than others. The maximum observed density is 6144.88, while the minimum density is 0.10, indicating large spatial variability in meteorite distribution (Kaiden, H, et al., 2006). Hotspots of meteorite activity can be identified through spatial cluster analysis, allowing researchers to pinpoint areas of increased meteorite impact rates and examine potential factors that contribute to these spatial patterns.

5.2 Temporal Trends:

As shown in Figure 2, temporal analysis shows interesting trends in meteorite landings over time (M. E., Zolensky ,2002). The time trend ranges from January 1, 1973, to January 22, 2501, indicating a comprehensive study of meteor impacts over the centuries. The density of meteorite impacts appears to have increased over time, suggesting a possible increase in meteorite activity in recent years or improvements in meteorite landing records. Further study of temporal changes and the identification of temporal clusters could provide valuable insights into the dynamic nature of meteoritic phenomena and their underlying mechanisms.

5.3 Classification and Spatial Distribution

Figure 3 shows the spatial distribution of meteorite classes ranging from c1/2-ung to Winconaite, Urellite-price, Urelite-an, Urelite, Iron, ungrouped, H3.3 and above (Weisberg M. K. et al., 2001). The spatial distribution of meteorite classes provides insights into the geological diversity of meteorite impacts and their formation. By classifying meteorites and studying their spatial distribution, researchers can elucidate the geological processes and parent bodies responsible for meteorite impacts, thereby improving our understanding of solar system dynamics and planetary evolution.

5.4 Implications for Planetary Science

The spatiotemporal analysis of meteor impacts performed in this study has important implications for planetary science and hazard assessment. By identifying spatial patterns, temporal trends, and meteorite classifications, researchers can assess the frequency, distribution, and characteristics of meteorite impacts globally (Goderis, S.,2012). This information is critical to understanding planetary dynamics, assessing the potential threat of meteor impacts to Earth and other celestial bodies, and developing planetary defence and risk reduction strategies.

6 Conclusion

In conclusion, the amalgamation of Geographic Information System (GIS) technology with meteorite impact data has yielded substantial insights into the spatial patterns, temporal trends, and geological diversity of meteorite occurrences on a global scale. The analysis of spatial data has brought to light the spatial variations in meteorite density, identifying concentrated areas of meteorite activity in specific geographical locations (Kerr,2001). These discoveries underscore the significance of spatial clustering analysis in identifying regions with heightened meteorite impact frequencies and comprehending the factors influencing spatial distribution. Moreover, the examination of temporal data has unveiled interesting patterns in meteorite landings over time, indicating possible shifts in meteorite activity or enhancements in recording techniques (Buddeet al., 2018). The observed rise in meteorite density over time emphasizes the dynamic nature of meteorite the spatial distribution of meteorite classes offers valuable insights into the geological diversity of meteorite impacts and the origins of meteoritic material, highlighting the importance of continuous research in this field. Categorizing meteorite classes and analysing their spatial distribution present opportunities for exploring the geological mechanisms and progenitor bodies accountable for meteorite impacts, thereby enhancing our comprehension of solar system dynamics. The significance of this investigation transcends to planetary science and risk assessment, underscoring the necessity to comprehend the frequency, distribution, and characteristics of meteorite impacts. Through the utilization of GIS technology and spatial analysis methodologies, scholars can further clarify the intricacies of meteorite occurrences, enriching our understanding of solar system evolution and planetary mechanisms.

7 References

Anders E. and Zinner E. (1993) Interstellar grains in primitive meteorites: Diamond, silicon carbide, and graphite. Meteoritics,28, 490–514.

Benedix G. K., McCoy T. J., Keil K., and Love S. G. (2000) A petrologic study of the IAB iron meteorites: Constraints on the formation of the IAB-winonaite parent body. Meteoritics & Planet. Sci., 35, 1127–1141.

Beckett J. R., Connolly H. C., and Ebel D. S. (2006) Chemical processes in calcium aluminium-rich inclusions: A mostly MAS view of melting and crystallization. In Meteorites and the Early Solar System II (D. S. Lauretta and H. Y. Mc-Sween Jr.,eds.), this volume. Univ. of Arizona, Tucson.

Bischoff A., Scott E. R. D., Metzler K., and Goodrich C. A. (2006) Nature and origins of meteoritic breccias. In Meteorites and the Early Solar System II (D. S. Lauretta and H. Y. McSween Jr., eds.), this volume. Univ. of Arizona, Tucson.

Burbine T. H., McCoy T. J., Meibom A., Gladman B., and KeilK. (2002) Meteoritic parent bodies: Their number and identi- fication. In Asteroids III (W. F. Bottke Jr. et al., eds.), pp. 653– 667. Univ. of Arizona, Tucson.

Buseck P. R. (1977) Pallasite meteorites mineralogy, petrology and geochemistry. Geochim. Cosmochim. Acta, 41, 711–740.

Debaille, V., Imae, N., Yamaguchi, A., Goderis, S., Mikouchi, T., Debouge, W., Hublet, G., Van Roosbroek, N.,Zekollari, H., Kojima, H. and Claeys. Ph. (2013): The 2012⊖2013 joint field campaign for collecting

Greshake A., Krot A. N., Meibom A., Weisberg M. K., and Keil K. (2002) Heavily-hydrated matrix lumps in the CH and metal- rich chondrites QUE 94411 and Hammadah al Hamra 237. Meteoritics & Planet. Sci., 37, 281–29 meteorites in Antarctica: an efficient collaboration between Japan and Belgium. Antarctic Meteorites XXXVI. Tokyo, 2013-11-14/15. National Institute of Polar Research, 11⊖12

Kaiden, H., Kojima, H. and Goderis, S. (2010): Collection of the Asuka 09 meteorites by the 51st Japanese Antarctic Research Expedition: a preliminary report. Antarctic Meteorites XXXIII. Tokyo, 2010-06-08/09. National Institute of Polar Research, 34⊖35.

Kallemeyn G. W., Boynton W. V., Willis J., and Wasson J. T.(1978) Formation of the Bencubbin polymict meteoritic breccia. Geochim. Cosmochim. Acta, 42, 507–515.

Kallemeyn G. W. and Wasson J. T. (1985) The compositional classification of chondrites: IV. Ungrouped chondritic meteorites and clasts. Geochim. Cosmochim. Acta, 49, 261–270.

Keil K. (1989) Enstatite meteorites and their parent bodies. Meteoritics, 24, 195–208.

Krot A. N., Keil K., Goodrich C. A., Scott E. R. D., and Weisberg M. K. (2004) Classification of Meteorites. In Treatise on Geochemistry, Vol. 1: Meteorites, Comets, and Planets (A. M. Davis, ed.), pp. 83–128. Elsevier, Oxford.

Lin Y. and Kimura M. (1998) Petrographic and mineralogical study of new EH melt rocks and a new enstatite chondrite grouplet. Meteoritics & Planet. Sci., 33, 501–511.

Lovering J. F. (1962) The evolution of the meteorites — Evidence for the co-existence of chondritic, achondritic and iron meteorites in a typical parent meteorite body. In Researches on Meteorites (C. B. Moore, ed.), pp. 179–197. Wiley, New York. Lovering J. F., Nichiporuk W., Chodos A., and Brown H. (1957).

McCoy T. J., Scott E. R. D., and Haack H. (1993) Genesis of the IIICD iron meteorites: Evidence from silicate-bearing inclusions. Meteoritics, 28, 552–560.

Mittlefehldt D. W. (2004) Achondrites. In Treatise on Geochemistry, Vol. 1: Meteorites, Comets, and Planets (A. M. Davis,ed.), pp. 291–324. Elsevier, Oxford.

Nagahara H. (1992) Yamato-8002: Partial melting residue on the unique chondrite parent body. Proc. NIPR Symp. Antarct. Meteorites, 5, 191–223.

Pourkhorsandi, H., Gattacceca, J., Rochette, P., D ' Orazio, M., Kamali, H., de Avillez, R., Letichevsky, S., Djamali, M., Mirnejad, H., Debaille, V. and Jull, A.J.T. (2019): Meteorites from the Lut Desert (Iran). Meteoritics & Planetary Science, 54, 1737⊖1763, doi: 10.1111/maps.13311.

Rochette, P., Sagnotti, L., Bourot-Denise, M., Consolmagno, G., Folco, L., Gattacceca, J., Osete, M.L. and Pesonen,L. (2003): Magnetic classification of stony meteorites: 1. Ordinary chondrites. Meteoritics & Planetary Science, 38, 251⊖268, doi: 10.1111/j.1945-5100. 2003.tb00263. x.

Rochette, P., Gattacceca, J., Bonal, L., Bourot-Denis, M., Chevrier, V., Clerc, J.P., Consolmagno, G., Folco, L., Gounelle, M., Kohout, T., Pesonen, L., Quirico, E., Sagnotti, L. and Skripnik, A. (2008): Magnetic classification of stony meteorites: 2. non-ordinary chondrites. Meteoritics & Planetary Science, 43, 959⊖980, doi: 10.1111/j.1945-5100. 2008.tb01092.

Rubin A. E. and Kallemeyn G. W. (1990) Lewis Cliff 85332: A unique carbonaceous chondrite. Meteoritics, 25, 215–225.

Rubin A. E., James J. A., Keck B. D., Weeks K. S., Sears D. W. G., and Jarosewich E. (1985) The Colony meteorite and variations in CO3 chondrite properties. Meteoritics, 20, 175–197.

Rubin A. E., Kallemeyn G. W., Wasson J. T., Clayton R. N.,Mayeda T. K., Grady M. M., and Verchovky A. B. (2001) Cujba: A new Bencubbin-like meteorite fall from Nigeria (abstract). In Lunar and Planetary Science XXXII, Abstract #1779. Lunar and Planetary Institute, Houston (CD-ROM).

Tsuchiya, N., Ishikawa, M., Satish-Kumar, M., Kawakami, T., Kojima, H., Kaiden, H., Miura, H., Suganuma, Y.,Abe, M., Sasaki, D., Chiba, M., Okada, Y., Hashizume, F., Grantham, G. and Goderis, S. (2012): Report ongeological, geomorphological and meteorite fieldwork in the Sør Rondane Mountains, Eastern DronningMaud Land, 2009⊖2010 (JARE-51). Nankyoku Shiryô (Antarctic Record), 56, 295⊖379 (in Japanese withEnglish abstract), doi: 10.15094/00009664.

Weisberg M. K. (2001) Sahara 00182, the first CR3 chondrite and formation of multi-layered chondrules (abstract). Meteoritics & Planet. Sci., 36, A222–A223.

Weisberg M. K., Prinz M., and Nehru C. E. (1990) The Bencubbin chondrite breccia and its relationship to CR chondrites and theALH85085 chondrite. Meteoritics, 25, 269–279.

Weisberg M. K., Prinz M., Clayton R. N., Mayeda T. K., Sugiura N., Zashu S., and Ebihara M. (2001) A new metal-rich chondrite grouplet. Meteoritics & Planet. Sci., 36, 401–418.

Zolensky M. E., Score R., Schutt J. W., Clayton R. N., and May-eda T. K. (1989) Lea County 001, an H5 chondrite and LeaCounty 002, an ungrouped chondrite. Meteoritics, 24, 227–232.

Zolensky M. E., Nakamura K., Gounelle M., Mikouchi T., Kasama T., Tachikawa O., and Tonui (2002) Mineralogy of Tagish Lake: An ungrouped type 2 carbonaceous chondrite. Meteoritics & Planet. Sci., 37, 737–761.